OMS Medical Evaluation of Functional Activities

Employee's Name			SSN (last 4 digits)	
Reason for OMS visit				
Occupational injury/illness		work assessment		
Personal injury/illness	Other (sp	pecify):		
Update/evaluation of status				
A. The employee is advise	d to resume reg	ular activities.		
B The employee is advise	d to limit activit	ies.		
The functional restrictions nee	eded from <i>(mo./day/yr.</i>)	_ through (mo./day/yr.)	
No climbing stairs or lade	ders	No twisting or bending	J	
No stooping or kneeling		No standing longer that	an ।	minutes per hour
No reaching above shoul	der	=	n	
No reaching below knee		=		·
No use of arm: left	, right	=	·	
		=		
	, right	= -	>	Journas
No fine manipulation		No operating motor ve		
Wear splint while working	g	Other, please describe	9:	
responsibility. If functional ressuitable duties are available e C. Insufficient information Proposed functional rest	is provided to exist the employee side where.	establish a medical ba Work absence proposed functional restricate a formal request for a re	e office responsible for persons sis for: extions and/or work absence	e, the supervisor should contact
Consult Employee Assista	ance Program	Immediately Next	available appointment	As planned on
Return to OMS for re-eva	luation on			
Negotiate leave with supe	ervisor			
Supervisor's name		Phone number	Supervisor contacted: Can accommodate Cannot accommod Unavailable; mess	
The OMS representative can be reached on 301-496-4411 to discuss the recommendations.	OMS representativ	ve	Date	Employee time in Employee
NIH 2558 (Rev. 4/04) Front			Distribution of copies:	Original — Supervisor Second — Employee Third — OMS files

Injury Treatment Instructions

Please read and follow the instructions marked below. Call the Occupational Medical Service (OMS) at 301-496-4411 if you have any questions regarding these instructions.

Sprain or Soft Tissue Injuries

- 1. Elevate the injured area to reduce swelling and pain.
- 2. Apply ice packs or cold compresses 10-20 minutes each hour while awake until pain subsides.
- Use the elastic bandage/splint on for _____ days. Remove and rewrap it as necessary. Be sure that it is snug, but not tight. Do not wear overnight unless otherwise instructed.
- Notify OMS if the injured area becomes cold or numb.

Cuts, Abrasions, or Burns

- 1. Keep the injured area clean, dry, and covered.
- 2. Change the bandage daily or if it becomes dirty or wet.
- 3. Call OMS if any of the following occur at the injury site: signs of infection (increasing redness, swelling, heat, pain, drainage) or excessive bleeding.
- 4. If the wound was sutured, the sutures will be removed in _____ days.

Fractures

- 1. Please see your physician as discussed for the fracture diagnosed by x-ray.
- Call your physician or seek urgent medical care at the nearest emergency room if any of the following occurs:

Burning, numbness, or discoloration of your fingers or toes.

Throbbing pain which doesn't decrease after elevating the injured limb on a pillow.

Back Injuries

- 1. Apply an ice pack for 10–20 minutes every hour until the pain subsides.
- 2. Sleep on a firm mattress either on your:

back with a pillow under your knees or side with a pillow between your knees.

Head Injuries

- 1. Take only the medications OMS has given to you for pain.
- 2. Do not take aspirin or other pain remedies.
- Someone should check you every 2 hours (including waking you from sleep) for 24 hours following the injury.
- 4. Call OMS or seek urgent medical care at the nearest emergency room if any of the following occurs:

Unusual irritability.

Definite changes in your behavior or personality.

Drowsiness or inability to be awakened.

Increasingly severe headache.

Dizziness or clumsy walking.

Slurred speech or changes in your vision or hearing.

Seizure or convulsion.

Arm or leg weakness.

Clear or bloody drainage from nose or ear.

Persistent vomiting.

Eye Injuries

- 1. Do not drive or operate power equipment if your eye is patched/covered.
- 2. If you need to remove your eye patch, use the provided clean supplies to replace the eye patch.
- 3. Call OMS immediately if eye pain increases or eye discharge occurs.

Non-Steroidal Anti-Inflammatory Medication (for example: aspirin, ibuprofen, naproxen)

- 1. Take the medication with food to decrease the risk of stomach irritation.
- Stop taking the medication and contact OMS if you notice any of the following: nausea, abdominal pain, diarrhea, vomiting, or black tarry stools.
- 3. Take the medicine at regular intervals:

Medication: _	
Schedule:	
ochedule	
Medication: _	
Schedule:	
Concadio.	